

GENERAL INSTRUCTIONS PRIOR TO SURGERY

DO NOT TAKE

THE FOLLOWING MEDICATIONS FOR TWO WEEKS BEFORE AND AFTER SURGERY (THEY PROMOTE BLEEDING)

Aclicin Act-3 Actiprofen	Decrin Dimetapp Spray Dipentum	Mobilis Morphalgin	Spraytish Sprein Strepfen Lozenges/Stepfen Surgam
Advil Aggrastat Aleve Anaprox Arcoxia Arthrexin Asacol Asasantin Aspalgia Aspirin Aspro Astrix	Disprin Dithranol Dolobid Drixine Ecotrin/Eliquis Entocort Feldene Fenac Fenpaed Fensaïd Fish Oil	Naprosyn Nurofen Junior Nurofen Naprogesic Nurolasts Orudis Oruvail Otrivin Omacar	Ticlopidine/Ticlid Toradil Tilcotil Tilodene Veganin Vincents Powders Vioxx Voltaren Warfarin Winsprin Capsules
Bex Brilinta Brufin	Glucosamine Hicin	Panafen Pentasa Persantin Pirox Ponstan Prexige Proxen Pradaxa Pyralin Rafen Rowasa	Xarelto <u>VITAMINS</u> Ginko Garlic Echinacea Vitamin E Feverfew/Flaxseed Oil/Fish Oil Ginger Ginseng St John's Wort
Candyl Canasa Cardiprin Cartia Celebrex Clinoril Clusinol Codiphen Colazal Chemgesic	Ibruprofen Indocid Indomed Inza Isocover Logicin Lyprinol	Salazopyrin Saldac Salofalk Salicylic Acid Cream Sinex Solcode Solprin	
Codox Codral Colazide Coplavix	Manuka Honey Marevan Maxigesic Mefic Mesasal Mobic		

IT IS SAFE

TO TAKE THE FOLLOWING IF YOU HAVE PAIN OR HEADACHE

Antibiotics Capadex Codalgin Diffiam Digesic	Doloxene Dymadon/Dimetapp Mersyndol Nasonex Painstop	Panadeine/Forte Panadol Panamax Panquil Paracetamol	Paradex Telfast/Claratyne/Zyrtec
----------------------------------------------------------	------------------------------------------------------------------	-----------------------------------------------------------------	-------------------------------------

If you develop a cold or gastroenteritis in the week before surgery, please contact the rooms on **9487.7877**.